

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/12

Paper 1 Pure Mathematics 1 (P1)

May/June 2017

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 75.

This document consists of **19** printed pages and **1** blank page.

- 1 (i) Find the coefficient of x in the expansion of $\left(2x - \frac{1}{x}\right)^5$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (ii) Hence find the coefficient of x in the expansion of $(1 + 3x^2)\left(2x - \frac{1}{x}\right)^5$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 2 The point A has coordinates $(-2, 6)$. The equation of the perpendicular bisector of the line AB is $2y = 3x + 5$.

(i) Find the equation of AB .

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find the coordinates of B .

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

3 (i) Prove the identity $\left(\frac{1}{\cos \theta} - \tan \theta\right)^2 \equiv \frac{1 - \sin \theta}{1 + \sin \theta}$. [3]

Cambridge Math & Physics
kcambridgemp.com

Kambridge Math & Physics
kambridgemp.com



- Cambridge Math & Prep
kamdbridgemp.com

Cambridge Math & Physics

kambridgemp.com


5 A curve has equation $y = 3 + \frac{12}{2-x}$.

(i) Find the equation of the tangent to the curve at the point where the curve crosses the x -axis. [5]

Cambridge Math & Physics

kambridgemp.com

Cambridge Math & Physics



The diagram shows the straight line $x + y = 5$ intersecting the curve $y = \frac{4}{x}$ at the points $A(1, 4)$ and $B(4, 1)$. Find, showing all necessary working, the volume obtained when the shaded region is rotated through 360° about the x -axis. [7]

Cambridge Math & Prep
kambridgemp.com

Kambridge Math & Physics
kambridgemp.com

- 7 (a) The first two terms of an arithmetic progression are 16 and 24. Find the least number of terms of the progression which must be taken for their sum to exceed 20 000. [4]

Cambridge Math & Physics

kambridgemp.com

- (b) A geometric progression has a first term of 6 and a sum to infinity of 18. A new geometric progression is formed by squaring each of the terms of the original progression. Find the sum to infinity of the new progression. [4]

Cambridge Math & Physics

kambridgemp.com

- 8** Relative to an origin O , the position vectors of three points A , B and C are given by

$$\overrightarrow{OA} = 3\mathbf{i} + p\mathbf{j} - 2p\mathbf{k}, \quad \overrightarrow{OB} = 6\mathbf{i} + (p+4)\mathbf{j} + 3\mathbf{k} \quad \text{and} \quad \overrightarrow{OC} = (p-1)\mathbf{i} + 2\mathbf{j} + q\mathbf{k},$$

where p and q are constants.

- (i)** In the case where $p = 2$, use a scalar product to find angle AOB .

[4]

Cambridge Math & Physics
kamdridgemp.com

Cambridge Math & Physics
kambriqemp.com

9 The equation of a curve is $y = 8\sqrt{x} - 2x$.

(i) Find the coordinates of the stationary point of the curve.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find an expression for $\frac{d^2y}{dx^2}$ and hence, or otherwise, determine the nature of the stationary point.

[2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (iii)** Find the values of x at which the line $y = 6$ meets the curve.

[3]

Cambridge Math & Physics
kambridgemp.com

- (iv)** State the set of values of k for which the line $y = k$ does not meet the curve.

[1]

[illegible]

10 The function f is defined by $f(x) = 3 \tan\left(\frac{1}{2}x\right) - 2$, for $-\frac{1}{2}\pi \leq x \leq \frac{1}{2}\pi$.

- (i) Solve the equation $f(x) + 4 = 0$, giving your answer correct to 1 decimal place. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (ii) Find an expression for $f^{-1}(x)$ and find the domain of f^{-1} . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Sketch, on the same diagram, the graphs of $y = f(x)$ and $y = f^{-1}(x)$.

[3]

Kambridge Math & Physics
kambridgemp.com

Kambridge Math & Physics
kambridgemp.com

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.