

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/33

Paper 3 Pure Mathematics 3 **(P3)**

May/June 2017

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 75.

This document consists of **19** printed pages and **1** blank page.

1 Prove the identity $\frac{\cot x - \tan x}{\cot x + \tan x} \equiv \cos 2x$.

[3]

Cambridge Math & Physics

kambridgeemp.com

- 2** Expand $(3 + 2x)^{-3}$ in ascending powers of x up to and including the term in x^2 , simplifying the coefficients. [4]

Cambridge Math & Physics

kambridgemp.com

- 3 Using the substitution $u = e^x$, solve the equation $4e^{-x} = 3e^x + 4$. Give your answer correct to 3 significant figures. [4]

Cambridge Math & Physics
kamdridgemp.com

- [4]

Cambridge Math & Physics

kambridgemp.com

- 5** A curve has equation $y = \frac{2}{3} \ln(1 + 3 \cos^2 x)$ for $0 \leq x \leq \frac{1}{2}\pi$.

(i) Express $\frac{dy}{dx}$ in terms of $\tan x$.

[4]

Cambridge Math & Physics

- (ii) Hence find the x -coordinate of the point on the curve where the gradient is -1 . Give your answer correct to 3 significant figures. [2]

Cambridge Math & Physics

kambridgemp.com


(i) Show by calculation that α is greater than 2.5. [2]

Cambridge Math & Physics
kamdbridgemp.com

(ii) Show that, if a sequence of values in the interval $0 < x < \pi$ given by the iterative formula $x_{n+1} = \pi + \tan^{-1}\left(\frac{1}{1 - x_n}\right)$ converges, then it converges to α . [2]

[illegible]

- (iii) Use this iterative formula to determine α correct to 3 decimal places. Give the result of each iteration to 5 decimal places. [3]



(i) Find the x -coordinate of M .

[4]

Cambridge Math & Physics
kamdridgemp.com

- (iii) The estimate found in part (ii) is denoted by E . Explain, without further calculation, whether another estimate found using the trapezium rule with four intervals would be greater than E or less than E . [1]

- 8** In a certain chemical reaction, a compound A is formed from a compound B . The masses of A and B at time t after the start of the reaction are x and y respectively and the sum of the masses is equal to 50 throughout the reaction. At any time the rate of increase of the mass of A is proportional to the mass of B at that time.

(i) Explain why $\frac{dx}{dt} = k(50 - x)$, where k is a constant. [1]

It is given that $x = 0$ when $t = 0$, and $x = 25$ when $t = 10$.

(ii) Solve the differential equation in part (i) and express x in terms of t . [8]

[illegible]

Kambridge Math & Physics
kambridgemp.com

9 Let $f(x) = \frac{3x^2 - 4}{x^2(3x + 2)}$.

(i) Express $f(x)$ in partial fractions.

[5]

Cambridge Math & Physics
kamdbridgemp.com

(ii) Hence show that $\int_1^2 f(x) \, dx = \ln\left(\frac{25}{8}\right) - 1$.

[5]

[illegible]

- 10** The points A and B have position vectors given by $\overrightarrow{OA} = \mathbf{i} - 2\mathbf{j} + 2\mathbf{k}$ and $\overrightarrow{OB} = 3\mathbf{i} + \mathbf{j} + \mathbf{k}$. The line l has equation $\mathbf{r} = 2\mathbf{i} + \mathbf{j} + m\mathbf{k} + \mu(\mathbf{i} - 2\mathbf{j} - 4\mathbf{k})$, where m is a constant.

(i) Given that the line l intersects the line passing through A and B , find the value of m . [5]

Cambridge Math & Physics
kamdridgemp.com

- (ii) Find the equation of the plane which is parallel to $\mathbf{i} - 2\mathbf{j} - 4\mathbf{k}$ and contains the points A and B .
Give your answer in the form $ax + by + cz = d$. [5]

Cambridge Math & Physics

kambridgemp.com

- $$z + (1 + i)w = i \quad \text{and} \quad (1 - i)z + iw = 1.$$

[6]

Cambridge Math & Physics
kambriqemp.com

- Cambridge Math & Physics
- kambridgemp.com

Kambridge Math & Physics
kambridgemp.com

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.