

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/32

Paper 3 Pure Mathematics 3 (P3)

October/November 2017

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.


At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 75.

This document consists of **19** printed pages and **1** blank page.

1



The diagram shows a sketch of the curve $y = \frac{3}{\sqrt{9-x^3}}$ for values of x from -1.2 to 1.2 .

- (i) Use the trapezium rule, with two intervals, to estimate the value of

$$\int_{-1.2}^{1.2} \frac{3}{\sqrt{9-x^3}} dx,$$

giving your answer correct to 2 decimal places.

[3]

.....

.....

.....

.....

.....

.....

.....

- (ii) Explain, with reference to the diagram, why the trapezium rule may be expected to give a good approximation to the true value of the integral in this case. [1]

.....

.....

.....

.....

.....

.....

.....

- 2** Showing all necessary working, solve the equation $2 \log_2 x = 3 + \log_2(x + 1)$, giving your answer correct to 3 significant figures. [5]

Cambridge Math & Physics
kambridgemp.com

- 3 By expressing the equation $\tan(\theta + 60^\circ) + \tan(\theta - 60^\circ) = \cot \theta$ in terms of $\tan \theta$ only, solve the equation for $0^\circ < \theta < 90^\circ$. [5]

Cambridge Math & Physics

kambridgemp.com

Kambridge Math & Physics
kambridgemp.com

- 4 The curve with equation $y = \frac{2 - \sin x}{\cos x}$ has one stationary point in the interval $-\frac{1}{2}\pi < x < \frac{1}{2}\pi$.

(i) Find the exact coordinates of this point.

[5]

Cambridge Math & Physics

kambridgemp.com

[2]

Cambridge Math & Physics

kanbridgemp.com

5 The variables x and y satisfy the differential equation

$$(x + 1) \frac{dy}{dx} = y(x + 2),$$

and it is given that $y = 2$ when $x = 1$. Solve the differential equation and obtain an expression for y in terms of x . [7]

[7]

Kambridge Math & Physics
kcambridgemp.com

Kambridge Math & Physics
kambridgemp.com

- 6** The equation of a curve is $x^3y - 3xy^3 = 2a^4$, where a is a non-zero constant.

(i) Show that $\frac{dy}{dx} = \frac{3x^2y - 3y^3}{9xy^2 - x^3}$. [4]

Cambridge Math & Physics

kambridgemp.com

- (ii) Hence show that there are only two points on the curve at which the tangent is parallel to the x -axis and find the coordinates of these points. [4]

Cambridge Math & Physics

kambridgemp.com

7 Throughout this question the use of a calculator is not permitted.

The complex number $1 - (\sqrt{3})i$ is denoted by u .

- (i) Find the modulus and argument of u .

[2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (ii) Show that $u^3 + 8 = 0$.

[2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (iii) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying both the inequalities $|z - u| \leq 2$ and $\operatorname{Re} z \geq 2$, where $\operatorname{Re} z$ denotes the real part of z . [4]

Kambridge Math & Physics
kambridgemp.com

8 Let $f(x) = \frac{8x^2 + 9x + 8}{(1-x)(2x+3)^2}$.

(i) Express $f(x)$ in partial fractions.

[5]

Cambridge Math & Physics
kamdbridgemp.com

- (ii) Hence obtain the expansion of $f(x)$ in ascending powers of x , up to and including the term in x^2 . [5]

Cambridge Math & Physics

kambridgemp.com

9 It is given that $\int_1^a x^{\frac{1}{2}} \ln x \, dx = 2$, where $a > 1$.

(i) Show that $a^{\frac{3}{2}} = \frac{7 + 2a^{\frac{3}{2}}}{3 \ln a}$. [5]

Kambridge Math & Physics
kambridgemp.com

(ii) Show by calculation that a lies between 2 and 4.

[2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Use the iterative formula

$$a_{n+1} = \left(\frac{7 + 2a_n^{\frac{2}{3}}}{3 \ln a_n} \right)^{\frac{2}{3}}$$

to determine a correct to 3 decimal places. Give the result of each iteration to 5 decimal places.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- Cambridge Math & Physics
kamdridgemp.com

- [illegible]

Kambridge Math & Physics
kambridgemp.com

Kambridge Math & Physics
kambridgemp.com

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.