



Cambridge International AS & A Level

CANDIDATE
NAME
CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS**9709/22**

Paper 2 Pure Mathematics 2

May/June 2025**1 hour 15 minutes**

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Any blank pages are indicated.

* 000080000002 *



Kambridge Math & Physics

kambridgemp.com





[3]

Kambridge Math & Physics

kambridgemp.com

[2]


[3]

[illegible]



Kambridge Math & Physics

kambridgemp.com



(a) Show that the x -coordinate of P satisfies the equation $x = -0.5 \ln(0.25 + 0.125 \sin 3x)$. [1]

Grade Math & Physics

(b) Use an iterative formula, based on the equation in part (a), to find the x -coordinate of P correct to 4 significant figures. Use an initial value of 0.5 and give the result of each iteration to 6 significant figures. [3]

[illegible]

Cambridge Math & Physics

kambridgeemp.com

5

$$p(x) = ax^4 + bx^3 + 13x^2 - 35x + 15,$$

where a and b are constants. It is given that $(2x - 1)$ and $(x - 3)$ are factors of $p(x)$.

(a) Find the values of a and b .

[4]

Kambridge Math & Physics



(b) Hence factorise $p(x)$.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Find the least positive value of θ in radians such that $p(\cot 2\theta) = 0$.

[2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN



Kambridge Math & Physics

kambridgemp.com

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN


$$12 \cos 2\phi \sin (2\phi + 30^\circ) = 5$$


[5]

Cambridge Math & Physics

kambridgeemp.com

Kambridge Math & Physics

kambridgemp.com



* 000080000015 *



Kambridge Math & Physics
kambridgemp.com





Kambridge Math & Physics
kambridgemp.com

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

