

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/33

Paper 3 Pure Mathematics 3 (P3)

May/June 2019

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 75.

This document consists of **20** printed pages.



- [4]

Cambridge Math & Physics

kambridgemp.com

2

[5]

Cambridge Math & Physics

kambridgemp.com

3 Let $f(\theta) = \frac{1 - \cos 2\theta + \sin 2\theta}{1 + \cos 2\theta + \sin 2\theta}$.

(i) Show that $f(\theta) = \tan \theta$.

[3]

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. A large, light gray watermark is oriented diagonally from the bottom-left towards the top-right. The watermark contains the text "Kambridge Math & Physics" in a bold sans-serif font, and below it, the website address "kambridgeemp.com" in a smaller, regular sans-serif font.

(ii) Hence show that $\int_{\frac{1}{6}\pi}^{\frac{1}{4}\pi} f(\theta) \, d\theta = \frac{1}{2} \ln \frac{3}{2}$.

[4]

Blank lined paper for writing.

- 4** The equation of a curve is $y = \frac{1 + e^{-x}}{1 - e^{-x}}$, for $x > 0$.

(i) Show that $\frac{dy}{dx}$ is always negative.

[3]

Cambridge Math & Physics

kambridgemp.com

- (ii)** The gradient of the curve is equal to -1 when $x = a$. Show that a satisfies the equation $e^{2a} - 4e^a + 1 = 0$. Hence find the exact value of a . [4]

Cambridge Math & Physics
kamdbridgemp.com


It is given that $y = 2$ when $x = 0$. Solve the differential equation obtaining an expression for y^2 in terms of x . [7]

Cambridge Math & Physics

kambridgemp.com

Kambridge Math & Physics
kambridgemp.com

6



The diagram shows the curve $y = x^4 - 2x^3 - 7x - 6$. The curve intersects the x -axis at the points $(a, 0)$ and $(b, 0)$, where $a < b$. It is given that b is an integer.

(i) Find the value of b .

[1]

.....

.....

.....

.....

.....

.....

.....

(ii) Hence show that a satisfies the equation $a = -\frac{1}{3}(2 + a^2 + a^3)$.

[4]

.....

.....

.....

.....

.....

.....

.....

.....

- Cambridge
kemp

7 The curve $y = \sin\left(x + \frac{1}{3}\pi\right) \cos x$ has two stationary points in the interval $0 \leq x \leq \pi$.

(i) Find $\frac{dy}{dx}$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) By considering the formula for $\cos(A + B)$, show that, at the stationary points on the curve, $\cos\left(2x + \frac{1}{3}\pi\right) = 0$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

[3]

Cambridge Math & Physics

kambridgemp.com

$$u = \frac{4i}{1 - (\sqrt{3})i}.$$

- (i) Express u in the form $x + iy$, where x and y are real and exact. [3]

Cambridge Math & Physics
kamdridgemp.com

- (ii) Find the exact modulus and argument of u .

[2]

.....

.....

.....

.....

.....

.....

.....

.....

- (iii) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z| < 2$ and $|z - u| < |z|$. [4]

Kambridge Math & Physics
kambridgemp.com

9 Let $f(x) = \frac{2x(5-x)}{(3+x)(1-x)^2}$.

(i) Express $f(x)$ in partial fractions.

[5]

Cambridge Math & Physics
kamdridgemp.com

- (ii) Hence obtain the expansion of $f(x)$ in ascending powers of x up to and including the term in x^3 . [5]

Cambridge Math & Physics

kambridgemp.com

- 10** The line l has equation $\mathbf{r} = \mathbf{i} + 2\mathbf{j} + 3\mathbf{k} + \mu(2\mathbf{i} - \mathbf{j} - 2\mathbf{k})$.

- (i) The point P has position vector $4\mathbf{i} + 2\mathbf{j} - 3\mathbf{k}$. Find the length of the perpendicular from P to l . [5]

Cambridge Math & Physics
kamdridgemp.com

- (ii) It is given that l lies in the plane with equation $ax + by + 2z = 13$, where a and b are constants. Find the values of a and b . [6]

Cambridge Math & Physics
kamdridgemp.com

Kambridge Math & Physics

kambridgemp.com

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.