

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/32

Paper 3 Pure Mathematics 3 **(P3)**

May/June 2017

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 75.

This document consists of **19** printed pages and **1** blank page.



- 1** Solve the equation $\ln(x^2 + 1) = 1 + 2 \ln x$, giving your answer correct to 3 significant figures. [3]

Cambridge Math & Physics

kambridgemp.com

2 Solve the inequality $|x - 3| < 3x - 4$.

[4]

Cambridge Math & Physics
cambridgeemp.com

- 3** (i) Express the equation $\cot \theta - 2 \tan \theta = \sin 2\theta$ in the form $a \cos^4 \theta + b \cos^2 \theta + c = 0$, where a , b and c are constants to be determined. [3]

Cambridge Math & Physics

kambridgemp.com

(ii) Hence solve the equation $\cot \theta - 2 \tan \theta = \sin 2\theta$ for $90^\circ < \theta < 180^\circ$.

[2]

Cambridge Math & Physics

kambridgemp.com

4 The parametric equations of a curve are

$$x = t^2 + 1, \quad y = 4t + \ln(2t - 1).$$

(i) Express $\frac{dy}{dx}$ in terms of t .

[3]

[illegible]

- (ii) Find the equation of the normal to the curve at the point where $t = 1$. Give your answer in the form $ax + by + c = 0$. [3]

Cambridge Math & Physics

kambridgemp.com

- $x = \frac{10}{(1+t)^2}$. At the beginning of the process $y = 100$.

[6]

Blank lined paper for writing.

- (ii) Find the exact value approached by the mass of B as t becomes large. State what happens to the mass of A as t becomes large. [2]

(i) It is given that u is a root of the equation $x^3 + ax^2 - 3x + b = 0$, where the constants a and b are real. Find the values of a and b . [4]

Cambridge Math & Physics
kamdridgemp.com

- (ii) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying both the inequalities $|z - u| < 1$ and $|z| < |z + i|$. [4]

Kambridge Math & Physics
kambridgemp.com

- 7 (i) Prove that if $y = \frac{1}{\cos \theta}$ then $\frac{dy}{d\theta} = \sec \theta \tan \theta$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (ii) Prove the identity $\frac{1 + \sin \theta}{1 - \sin \theta} \equiv 2 \sec^2 \theta + 2 \sec \theta \tan \theta - 1$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (iii) Hence find the exact value of $\int_0^{\frac{1}{4}\pi} \frac{1 + \sin \theta}{1 - \sin \theta} d\theta$. [4]

8 Let $f(x) = \frac{5x^2 - 7x + 4}{(3x + 2)(x^2 + 5)}$.

(i) Express $f(x)$ in partial fractions.

[5]

Cambridge Math & Physics
kamdbridgemp.com

- (ii) Hence obtain the expansion of $f(x)$ in ascending powers of x , up to and including the term in x^2 . [5]

Cambridge Math & Physics
kcambridge.com

- (i) Find the position vector of the foot of the perpendicular from A to l . Hence find the position vector of the reflection of A in l . [5]

[illegible]


- (ii) Find the equation of the plane through the origin which contains l . Give your answer in the form $ax + by + cz = d$. [3]

Cambridge Math & Physics
cambridgep.com

- (iii) Find the exact value of the perpendicular distance of A from this plane. [3]

[illegible]

10



The diagram shows the curve $y = x^2 \cos 2x$ for $0 \leq x \leq \frac{1}{4}\pi$. The curve has a maximum point at M where $x = p$.

- (i) Show that p satisfies the equation $p = \frac{1}{2} \tan^{-1} \left(\frac{1}{p} \right)$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

- (ii) Use the iterative formula $p_{n+1} = \frac{1}{2} \tan^{-1} \left(\frac{1}{p_n} \right)$ to determine the value of p correct to 2 decimal places. Give the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

- (iii) Find, showing all necessary working, the exact area of the region bounded by the curve and the x -axis. [5]

Cambridge Math & Physics

kambridgemp.com

Kambridge Math & Physics
kambridgemp.com

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.