

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/32

Paper 3 Pure Mathematics 3 (P3)

October/November 2018

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 75.

This document consists of **19** printed pages and **1** blank page.



- [4]

Kambridge Math & Physics

- 2** Showing all necessary working, solve the equation $\sin(\theta - 30^\circ) + \cos \theta = 2 \sin \theta$, for $0^\circ < \theta < 180^\circ$. [4]

Cambridge Math & Physics
kambridgemp.com

- 3 (i) Find $\int \frac{\ln x}{x^3} dx$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (ii) Hence show that $\int_1^2 \frac{\ln x}{x^3} dx = \frac{1}{16}(3 - \ln 4)$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4 Showing all necessary working, solve the equation

$$\frac{e^x + e^{-x}}{e^x + 1} = 4,$$

giving your answer correct to 3 decimal places.

[5]

Kambridge Math & Physics

- 5** The equation of a curve is $y = x \ln(8 - x)$. The gradient of the curve is equal to 1 at only one point, when $x = a$.

(ii) Show that a satisfies the equation $x = 8 - \frac{8}{\ln(8-x)}$. [3]

Cambridge Math & Physics

kambridgemp.com

- (ii) Verify by calculation that a lies between 2.9 and 3.1.

[2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (iii) Use an iterative formula based on the equation in part (i) to determine a correct to 2 decimal places. Give the result of each iteration to 4 decimal places.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- Cambridge Math & Physics
kamdridgemp.com

Kambridge Math & Physics
kambridgemp.com

7 A curve has equation $y = \frac{3 \cos x}{2 + \sin x}$, for $-\frac{1}{2}\pi \leq x \leq \frac{1}{2}\pi$.

(i) Find the exact coordinates of the stationary point of the curve.

[6]

Cambridge Math & Physics

kambridgemp.com

- (ii) The constant a is such that $\int_0^a \frac{3 \cos x}{2 + \sin x} \, dx = 1$. Find the value of a , giving your answer correct to 3 significant figures. [4]

Cambridge Math & Physics

kambridgemp.com

8 Let $f(x) = \frac{7x^2 - 15x + 8}{(1 - 2x)(2 - x)^2}$.

(i) Express $f(x)$ in partial fractions.

[5]

Cambridge Math & Physics
kamdbridgemp.com

- (ii) Hence obtain the expansion of $f(x)$ in ascending powers of x , up to and including the term in x^2 . [5]

Cambridge Math & Physics

kambridgemp.com

- 9 (a) (i) Without using a calculator, express the complex number $\frac{2+6i}{1-2i}$ in the form $x+iy$, where x and y are real. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (ii) Hence, without using a calculator, express $\frac{2+6i}{1-2i}$ in the form $r(\cos \theta + i \sin \theta)$, where $r > 0$ and $-\pi < \theta \leq \pi$, giving the exact values of r and θ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- [5]

Cambridge
kamdbridgemp.com

(i) Find the position vector of A .

[3]

Cambridge Math & Physics

kambridgemp.com

- (ii) Calculate the acute angle between l and p .

[4]

Cambridge Math & Physics

kambridgemp.com

[Question 10 (iii) is printed on the next page.]

[4]

Cambridge Math & Physics

kanbridgemp.com

Cambridge Math & Physics

kambridgemp.com

Kambridge Math & Physics
kambridgemp.com

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.