



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/12

Paper 1 Pure Mathematics 1

October/November 2025

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Any blank pages are indicated.



- 1 (a) Express $9x^2 - 36x + 8$ in the form $p(x+q)^2 + r$, where p , q and r are constants. [2]

[illegible]

- (b) Hence find the set of values of the constant k for which the equation $9x^2 - 36x + 8 = k$ has no real roots. [1]

Grade Math & Phys

- (c) Find the exact roots of the equation $9x^2 - 36x + 8 = -15$. [2]

[illegible]



- 2** Find the term independent of x in the expansion of $\left(2x^2 - \frac{3}{x}\right)^6$. [3]

Cambridge Math & Physics

cambridgeemp.com





- 3 (a) The graph of $y = f(x)$ is transformed to the graph of $y = f(3x) + 2$.

Describe fully the two transformations which have been combined to give the resulting graph. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) A different graph has equation $y = g(x)$. This graph is stretched by scale factor 3 in the y -direction and then reflected in the y -axis.

Write down the equation of the transformed graph in terms of the function g . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....





- 4 The equation of a curve is such that $\frac{dy}{dx} = kx^3 + \frac{2}{x^2}$, where k is a constant. The curve passes through the point $S(2, 20)$ and the gradient of the curve at S is $\frac{65}{2}$.

(a) Find the value of k .

[1]


- (b)** The coordinates of a point T on the curve are $(1, t)$.

Find the value of t .

[5]



5



The equation of a curve is $y = 4x^{\frac{1}{2}} - x$. The curve has a maximum point when $x = a$ and crosses the x -axis at the point with coordinates $(b, 0)$, where $b > 0$. The shaded region is bounded by the curve, the line $x = a$ and the x -axis (see diagram).

- (a)** Find the value of a .

[3]

[illegible]



(b) Find the exact area of the shaded region.


[5]

Cambridge Math & Physics



6 (a) Sketch the graph of $y = 3 \sin x + 2$ for $0 \leq x \leq 2\pi$.

[2]



(b) Determine the number of solutions in the interval $0 \leq x \leq 2\pi$ of each of the following equations.

(i) $3 \sin x + 2 = x$

[1]

.....

.....

.....

.....

.....

(ii) $3 \sin x + 2 = 5 - x$

[1]

.....

.....

.....

.....

.....





(c) Solve the equation $3 \sin x + 2 = 5 \cos^2 x - 1$ for $0 \leq x \leq 2\pi$.

[5]

Cambridge Math & Physics

kambridgemp.com





- 7 The coordinates of the points P and Q are $(1, 1)$ and $(7, 11)$ respectively. The line segment PQ forms a diameter of a circle.

(a) Find the equation of the circle.

[4]

Cambridge Math & Physics
kambriidgemp.com

(b) Find the equation of the tangent to the circle at the point Q .

[3]

[illegible]



- (c) The other point on the circle with x -coordinate 7 is R .

Find the coordinates of the point of intersection of the tangent at Q with the tangent at R . [4]

Cambridge Math & Physics

kambridgeemp.com



- 8 The first three terms of a geometric progression are a , b and c respectively, where a , b and c are positive constants. The first three terms of an arithmetic progression are a , b and $-3c$ respectively.

(a) Show that $a^2 - 10ac + 9c^2 = 0$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

It is now given that $a = 9$ and c takes the smaller of its two possible values.

(b) (i) Find the sum to infinity of the geometric progression. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....





.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find the sum of the first 20 terms of the arithmetic progression. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Kambridge Math & Physics

kambridgemp.com





9 The function f is defined by $f(x) = \frac{4}{(3x-6)^2} + \frac{1}{(3x-6)^3}$ for $x > 2$.

- (a) Find an expression for $f'(x)$ and hence determine whether f is an increasing function, a decreasing function or neither. [4]

Cambridge Math & Physics
kambriqemp.com

- (b) State whether f^{-1} exists. Give a reason for your answer. [1]

[illegible]



The function g is defined by $g(x) = 4x - 3$ for $x > a$.

- (c) Find the range of g in terms of the constant a . [1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (d) Find the set of values of a for which the composite function fg exists. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....





The diagram shows a circle with centre A and radius r passing through points B , C and D . A larger circle of radius s has centre C and passes through B and D . The length BD is also s .

(a) Show that $s = \sqrt{3}r$.

[2]





- (b)** Find an expression for the area of the shaded region. Give your answer in the form $(a + b\pi)r^2$, where a and b are constants to be found. [7]

Cambridge Math & Physics

kambridgemp.com





Kambridge Math & Physics

kambridgemp.com



Kambridge Math & Physics
kambridgemp.com





Kambridge Math & Physics
kambridgemp.com

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

