

Cambridge International AS & A Level

MATHEMATICS**9709/15**

Paper 1 Pure Mathematics 1

May/June 2025

MARK SCHEME

Maximum Mark: 75

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2025 series for most Cambridge IGCSE, Cambridge International A and AS Level components, and some Cambridge O Level components.

This document consists of **24** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptions for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Mathematics-Specific Marking Principles

- 1 Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
- 2 Unless specified in the question, non-integer answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
- 3 Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
- 4 Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
- 5 Where a candidate has misread a number or sign in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 A or B mark for the misread.
- 6 Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

Kambridge Math & Physics
kambridgemp.com










Annotations guidance for centres


Examiners use a system of annotations as a shorthand for communicating their marking decisions to one another. Examiners are trained during the standardisation process on how and when to use annotations. The purpose of annotations is to inform the standardisation and monitoring processes and guide the supervising examiners when they are checking the work of examiners within their team. The meaning of annotations and how they are used is specific to each component and is understood by all examiners who mark the component.


We publish annotations in our mark schemes to help centres understand the annotations they may see on copies of scripts. Note that there may not be a direct correlation between the number of annotations on a script and the mark awarded. Similarly, the use of an annotation may not be an indication of the quality of the response.

The annotations listed below were available to examiners marking this component in this series.

Annotations

Annotation	Meaning
	More information required
	Accuracy mark awarded zero
	Accuracy mark awarded one
	Independent accuracy mark awarded zero
	Independent accuracy mark awarded one
	Independent accuracy mark awarded two
	Benefit of the doubt
	Blank Page
	Incorrect
Dep	Used to indicate DM0 or DM1

Annotation	Meaning
DM1	Dependent on the previous M1 mark(s)
FT	Follow through
	Indicate working that is right or wrong
Highlighter	Highlight a key point in the working
ISW	Ignore subsequent work
J	Judgement
JU	Judgement
M0	Method mark awarded zero
M1	Method mark awarded one
M2	Method mark awarded two
MR	Misread
O	Omission or Other solution
Off-page comment	Allows comments to be entered at the bottom of the RM marking window and then displayed when the associated question item is navigated to.
On-page comment	Allows comments to be entered in speech bubbles on the candidate response.
PE	Judgment made by the PE
Pre	Premature approximation
SC	Special case
SEEN	Indicates that work/page has been seen

Annotation	Meaning
SF	Error in number of significant figures
	Correct
TE	Transcription error
XP	Correct answer from incorrect working

Kambridge Math & Physics
kambridgemp.com

PUBLISHED**Mark Scheme Notes**

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

- M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B** Mark for a correct result or statement independent of method marks.
- DM or DB** When a part of a question has two or more ‘method’ steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- FT** Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.
- A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT above).
 - For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures or would be correct to 3 significant figures if rounded (1 decimal place for angles in degrees).
 - The total number of marks available for each question is shown at the bottom of the Marks column.
 - Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
 - Square brackets [] around text or numbers show extra information not needed for the mark to be awarded.

Abbreviations

AEF/OE	Any Equivalent Form (of answer is equally acceptable) / Or Equivalent
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
CAO	Correct Answer Only (emphasising that no ‘follow through’ from a previous error is allowed)
CWO	Correct Working Only
ISW	Ignore Subsequent Working
SOI	Seen Or Implied
SC	Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)
WWW	Without Wrong Working
AWRT	Answer Which Rounds To

PUBLISHED

Question	Answer	Marks	Guidance
1	$[y =] \left\{ \frac{12}{3 \times 2} (2x - 5)^3 \right\} \left\{ + \frac{8}{2} x^2 \right\} [+c]$	B1 B1	OE Terms may be unsimplified. May see $[y =] \{16x^3\} \{-116x^2\} \{+300x\} [+c]$.
	$4 = 2 \times (2 \times 2 - 5)^3 + 4 \times 2^2 + c \Rightarrow c = -10]$	M1	Sub (2, 4) correctly into <i>their</i> integrated expression with a '+ c'.
	y or f or $f(x) = 2(2x - 5)^3 + 4x^2 - 10$	A1	May see $y = 16x^3 - 116x^2 + 300x - 260$. 'y = ' or 'f(x) = ' or 'f = ' can be implied if seen in working.
		4	

PUBLISHED

Question	Answer	Marks	Guidance
2	x^2 coeff = $10 \times 3^3 \times a^2$ and $6 \times 6^2 \times (-1)^2$ allow correct terms with x^2	B1	Expect $270a^2$ and 216. Terms may be seen separately. Allow if seen in an expansion. Combinations must be evaluated.
	x coeff = $5 \times 3^4 \times a$ and $4 \times 6^3 \times (-1)$ allow correct terms with x	B1	Expect $405a$ and -864 . Terms may be seen separately. Allow if seen in an expansion. Combinations must be evaluated.
	$270a^2 + 216 = 6(405a - 864)$	M1	OE For forming a correct quadratic equation using <i>their</i> 4 terms only.
	$45a^2 - 405a + 900 = 0 \Rightarrow 45(a - 4)(a - 5) = 0$	DM1	For evidence of a correct method of solving. If quadratic formula used a full substitution must be seen.
	$a = 4$ or $a = 5$	A1	Only dependent on the first M1.
			5

PUBLISHED

Question	Answer	Marks	Guidance
3(a)	$4\left(x - \frac{1}{2}\right)^2 [-2=0]$ or $(2x-1)^2 [-2=0]$ or $\left(x - \frac{1}{2}\right)^2 \left[-\frac{1}{2}=0\right]$	M1	OE Must deal with the coefficient of x^2 and x correctly to produce an $(ax + b)^2$ term.
	$x = \frac{1}{2} \pm \sqrt{\frac{1}{2}}$	A1	OE, e.g. $x = \frac{1}{2}(1 \pm \sqrt{2})$. SC B1 only for correct solutions from another method.
		2	
3(b)	$\tan \theta = \frac{1}{2}(1 \pm \sqrt{2})$	M1	Setting $\tan \theta = \textit{their } x$ for at least one value of <i>their</i> x . May restart and solve the quadratic in $\tan \theta$.
	$[\theta =] 50.4, 168.3$ AWRT and no other answers in the range $0 < \theta < 180$	A1 A1	SC A1 only for AWRT 0.879 and 2.94 radians. SC M0 B1 B1 for answers only. SC M0 B1 (only) for both answers only in radians.
		3	

PUBLISHED

Question	Answer	Marks	Guidance
4	Intersects x -axis when $x^4 = 1 \Rightarrow x = 1$	B1	WWW May be seen without working. May be seen as a limit in the integration. Allow ± 1 if $x = 1$ is seen as the lower limit in the integration.
	$V = [\pi] \int y^2 dx = [\pi] \int \left(x^2 - \frac{1}{x^2}\right)^2 dx = [\pi] \int \left(x^4 - 2 + \frac{1}{x^4}\right) dx$ with attempt at integration	*M1	With some attempt at squaring (accept $x^4 \pm \frac{1}{x^4}$ as evidence).
	$\frac{1}{5}x^5 - 2x - \frac{1}{3x^3}$	A1	
	$[\pi] \left\{ \left(\frac{1}{5} \times 2^5 - 2 \times 2 - \frac{1}{3 \times 2^3} \right) - \left(\frac{1}{5} - 2 - \frac{1}{3} \right) \right\}$	DM1	Use of limits, $x = 2$ and <i>their</i> $x = 1$, min evidence if correct answer: $\frac{283}{120} + \frac{32}{15}$. If answer incorrect, substitution of limits must be clear. DM0 for use of the limit $x = 0$.
	$\pi \times 4.49\dots = 14.11$	A1	AWRT, WWW Allow $\frac{539}{120}\pi$ or 14.1, dependent on the first M1 and A1.
		5	

PUBLISHED

Question	Answer	Marks	Guidance
5(a)	Angle $CAD = \frac{\pi}{3}$ or 60° , $CD = \sqrt{75}$ or $5\sqrt{3}$	B1	OE For either $\hat{C}AD$ or CD . Allow $\hat{C}AD = 1.05$, $CD = 8.7$ or 8.66 .
	[Perimeter =] $\left\{10 \times \frac{\pi}{3}\right\} \{+\sqrt{75} + 5\}$ or $\left\{\frac{60}{360} \times 2\pi(10)\right\} \{+\sqrt{75} + 5\}$	B1 B1	OE B1 for the two sides and B1 for arc length (allow 10.5).
	= 24.1 cm	B1	
		4	
5(b)	Area = $\frac{1}{2} \times 10^2 \times \frac{\pi}{3} - \frac{1}{2} \times 5 \times \sqrt{75}$ or $\frac{60}{360} \times \pi \times 10^2 - \frac{1}{2} \times 5 \times \sqrt{75}$	M1	OE Use of sector area formula minus triangle area formulae, with <i>their</i> angle CAD and <i>their</i> side CD .
	= 30.7 cm ²	A1	
		2	

PUBLISHED

Question	Answer	Marks	Guidance
6(a)	Eleventh birthday: Father $10+10\times 5 [= 60]$	M1	For correct use of AP formula OE.
	Mother $10\times 1.2^{10} [= 61.9(2)]$	M1	For correct use of GP formula OE.
	60 and 61.9	A1	Both answers. Accept 2 sf answers.
		3	
6(b)	Father $\frac{18}{2}(2\times 10+17\times 5) = 945$	M1A1	A1 may be implied by a correct final answer.
	Mother $\frac{10(1-1.2^{18})}{(1-1.2)} = 1281.17$	M1A1	A1 may be implied by a correct final answer.
	Total = 2226.17	A1	Accept 3 or more sf answers. Ignore $S_{17} (= 1909.3)$ as an extra answer.
		5	

Question	Answer	Marks	Guidance
7(a)	$\frac{7-p}{3-6} = -\frac{2}{3}$ or use of straight line equations with $m = -\frac{2}{3}$, $(x, y) = (3, 7)$ with $(6, p)$ substituted.	M1	OE Expect $y - 7 = \left(-\frac{2}{3}\right)(x - 3)$ with $(6, p)$ substituted or $y = \frac{-2}{3}x + 9$ with $(6, p)$ substituted.
	$p = 5$	A1	
		2	
7(b)	$(\{4\}, \{their\ 2p-7\})$	B1	WWW
		B1FT	Extra solutions lose both marks.
		2	
7(c)	Mid-point AB is $(4.5, 6)$	B1	
	Gradient of perp bisector $\left[\begin{array}{l} = \frac{-1}{-\frac{2}{3}} \\ = \frac{3}{2} \end{array} \right]$	B1	
	Equation $y - 6 = \frac{3}{2}(x - 4.5)$	M1	OE. Must be using <i>their</i> mid-point and <i>their</i> perpendicular gradient.
	Crosses axes at $(0, -0.75), (0.5, 0)$	DM1	Correct use of <i>their</i> perpendicular bisector equation to find the x - and y -intercept.
	Area = 0.1875 (accept 3 sf accuracy)	A1	OE, e.g. $\frac{3}{16}$. Last 2 marks can be gained by integrating the line equation between zero and 0.5.
		5	

PUBLISHED


Question	Answer	Marks	Guidance
8(a)	$9 = 1 + a + b + 5$	B1	
	$\frac{dy}{dx} = 3x^2 + 2ax + b$	B1	
	Gradient = 0 at (1, 9) so $0 = 3 + 2a + b$	M1	Setting <i>their</i> $\frac{dy}{dx}$ to zero and substituting $x = 1$.
	Attempt to solve <i>their</i> linear equations simultaneously	DM1	Can be implied by their answers.
	$a = -6, b = 9$	A1	WWW
		5	
8(b)	$\frac{dy}{dx} = 3x^2 - 12x + 9 = 0$	M1	Setting <i>their</i> $\frac{dy}{dx}$ to zero.
	Solution [leading to $x = 1$ or $x = 3$]	DM1	Solving <i>their</i> 3-term quadratic
	(3, 5) or $x = 3, y = 5$	A1	WWW Ignore (1, 9) if given as a second answer. Only dependent on the first M1.
		3	

PUBLISHED

Question	Answer	Marks	Guidance
8(c)	At $x = 5$, $\frac{dy}{dx} = 3 \times 5^2 - 12 \times 5 + 9$	M1	Substituting $x = 5$ into <i>their</i> $\frac{dy}{dx}$. May be implied.
	$6 = \textit{their} 24 \times \frac{dx}{dt}$ or $\textit{their} 24 = 6 \times \frac{dt}{dx}$	M1	OE Use of chain rule SOI $\left(\frac{dy}{dt} = \frac{dy}{dx} \times \frac{dx}{dt}\right)$. Linking correctly $\frac{dx}{dt}$ (or $\frac{dt}{dx}$), <i>their</i> 24 and 6.
	$\frac{dx}{dt} = \frac{1}{4}$	A1	OE
		3	

PUBLISHED

Question	Answer	Marks	Guidance
9(a)	{Stretch} {factor 3} {in y-direction}	B2,1,0	If 2 or more stretches or extra transformations, give B0 for stretches.
	{Translation or shift} $\begin{pmatrix} \{\pi\} \\ \{2\} \end{pmatrix}$	B2,1,0	Translation may be split to $\begin{pmatrix} \pi \\ 0 \end{pmatrix}$ before the stretch and $\begin{pmatrix} 0 \\ 2 \end{pmatrix}$ after the stretch. If both vectors correct but ‘translation’/’shift’ not stated, then B1 only for the translations.
	Alternative Method for Question 9(a)		
	{Translation or shift} $\begin{pmatrix} \{\pi\} \\ \left\{\frac{2}{3}\right\} \end{pmatrix}$ or $\begin{pmatrix} \pi \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ \frac{2}{3} \end{pmatrix}$	B2,1,0	If two or more stretches or any extra incorrect transformation is given, then B0 for the stretches. If order incorrect, maximum of 3/4.
	followed by {stretch} {factor 3} {in y direction}	B2,1,0	
		4	

Question	Answer	Marks	Guidance
<p>9(b)</p> 		B1 †	Graph of $f(x)$ with correct domain.
		B1 †	For $g(x)$ being a decreasing function in the domain π to 2π .
		B1	Domain π to 2π for $g(x)$.
		B1	Range for g approximately correct (should be from 5 to -1).
<p>9(c)</p>	$[g^{-1}] f\left(\frac{\pi}{3}\right) = [g^{-1}]\left(\frac{1}{2}\right)$	B1	May be implied by correct substitution later.
	$[g^{-1}(x) \text{ or } y =] \cos^{-1}\left(\frac{x-2}{3}\right) + \pi \text{ or } [x =] \cos^{-1}\left(\frac{y-2}{3}\right) + \pi$	M1	Finding inverse of g . Allow one sign error. Alt: $3 \cos(x - \pi) + 2 = \frac{1}{2}$
	$[g^{-1}\left(\frac{1}{2}\right) =] \cos^{-1}\left(\frac{\frac{1}{2}-2}{3}\right) + \pi$	DM1	Substituting <i>their</i> $x = \frac{1}{2}$ into <i>their</i> expression for $g^{-1}(x)$. Alt: Use of arccos for $\cos(x - \pi) = -\frac{1}{2} \Rightarrow x - \pi = \frac{2\pi}{3}$
	$[g^{-1}(x) =] \frac{2\pi}{3} + \pi = \frac{5\pi}{3} \text{ or } 5.24 \text{ AWRT}$	A1	
		4	

PUBLISHED

Question	Answer	Marks	Guidance
9(d)	The domain of f does not include the whole of the range of g	B1	OE, but must mention range of g and domain of f.
Alternative Method for Question 9(d)			
	Show clearly that a particular value or set of values in the domain of g gives a value of $g(x)$ which is outside the domain of f	B1	Value of x for substitution into $g(x)$ must be in the ranges: $\pi \leq x < 4.322$ or $5.447 < x \leq 2\pi$.
		1	

Kambridge Math & Physics

kambridgemp.com

PUBLISHED

Question	Answer	Marks	Guidance
10(a)	Centre is $(-2, 4)$	B1	
	Gradient of perpendicular = $\frac{8-4}{2-(-2)}$ [$= 1$]	M1	OE, e.g. using equation of the radius.
	Equation of tangent is $y - 8 = -1(x - 2)$	M1	Using a correct point and <i>their</i> tangent gradient from an attempt to find $\frac{dy}{dx}$, or the negative reciprocal of <i>their</i> perpendicular gradient.
	$x + y - 10 = 0$ or $-x - y + 10 = 0$	A1	CAO – this form is required. Answer without working, allow maximum of 2/4.
Alternative Method for Question 10(a)			
	$y = [\pm](28 - x^2 - 4x)^{\frac{1}{2}} + 4$ or $y = [\pm](32 - (x+2)^2)^{\frac{1}{2}} + 4$	B1	OE; a correct expression for y .
	$\frac{dy}{dx} = [\pm]\frac{1}{2}(28 - x^2 - 4x)^{-\frac{1}{2}}(-2x - 4)$ [$= -1$ at $x = 2$]	M1	OE Differentiating y with no more than one sign error.
	Equation of tangent is $y - 8 = -1(x - 2)$	M1	Using a correct point and <i>their</i> tangent gradient from an attempt to find $\frac{dy}{dx}$, or the negative reciprocal of <i>their</i> perpendicular gradient.
	$x + y - 10 = 0$ or $-x - y + 10 = 0$	A1	CAO – this form is required. Answer without working, allow maximum of 2/4.

PUBLISHED

Question	Answer	Marks	Guidance
	Alternative Method 2 for Question 10(a)		
	$2x + 2y \frac{dy}{dx} + 4 - 8 \frac{dy}{dx} = 0$	B1	Differentiating implicitly.
	$\frac{dy}{dx} = \frac{(2x+4)}{(8-2y)}$ [$= -1$ at $(2, 8)$]	M1	Rearranging to make $\frac{dy}{dx}$ the subject, May substitute $(2, 8)$ and then rearrange to find $\frac{dy}{dx}$.
	Equation of tangent is $y - 8 = -1(x - 2)$	M1	Using a correct point and <i>their</i> tangent gradient from an attempt to find $\frac{dy}{dx}$, or the negative reciprocal of <i>their</i> perpendicular gradient.
	$x + y - 10 = 0$ or $-x - y + 10 = 0$	A1	CAO – this form is required. Answer without working, allow maximum of 2/4.
		4	

PUBLISHED

Question	Answer	Marks	Guidance
10(b)	$(k - 3y)^2 + y^2 + 4(k - 3y) - 8y - 12 = 0$ or $(k - 3y + 2)^2 + (y - 4)^2 = 32$	M1*	Sub $x = k - 3y$ or $y = \frac{k - x}{3}$ into circle equation. $y = \frac{k - x}{3}$ gives: $\frac{(k - x)^2}{9} - 8\frac{k - x}{3} + x^2 + 4x - 12 = 0$ or $(x + 2)^2 + \left(\frac{k - x}{3} - 4\right)^2 = 32.$
	$9y^2 - 6ky - 20y + y^2 + k^2 + 4k - 12 = 0$	DM1	OE All squared brackets expanded to give a quadratic equation in y (or x , which gives: $9x^2 + 60x - 2kx + k^2 + x^2 - 24k - 108 = 0$ or $\frac{x^2 + k^2 - 2kx}{9} - \frac{8k - 8x}{3} + 16 + x^2 + 4x + 4 = 32.$)
	$b^2 - 4ac = (6k + 20)^2 - 4 \times 10 \times (k^2 + 4k - 12)$	M1**	OE Factorising out y (or x) from <i>their</i> quadratic (this is the quadratic where the previous DM1 or possibly DM0 was awarded) to identify ' b ' and then finding the discriminant. From equation in x , this is: $(60 - 2k)^2 - (4)(10)(k^2 - 24k - 108).$
	$-4k^2 + 80k + 880 < 0$	DM1	OE Simplify the discriminant and setting it to less than zero. From eliminating y this is: $-36k^2 + 720k + 7920 < 0$ Only dependent on the previous M1.
	$k^2 - 20k - 220 > 0$	A1	AG WWW

PUBLISHED

Question	Answer	Marks	Guidance
10(b)	Alternative Method for Question 10(b)		
	Centre of circle $(-2, 4)$ and radius of circle is $\sqrt{32}$	B1	
	Distance of $x + 3y - k$ from $(-2, 4) = \frac{1(-2) + 3(4) - k}{\sqrt{1^2 + 3^2}}$	M1*	Correct use of ‘distance of a line from a point formula’ with <i>their</i> centre coordinates.
	Setting $\frac{1(-2) + 3(4) - k}{\sqrt{1^2 + 3^2}} > \sqrt{32}$	DM1	Setting distance to be greater than <i>their</i> radius.
	Squaring both sides	DM1	Removing the square roots.
	Rearranging to $k^2 - 20k - 220 > 0$	A1	AG
		5	