



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/13

Paper 1 Pure Mathematics 1

May/June 2021

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Any blank pages are indicated.



Kambridge Math & Physics
kambridgemp.com

- 1** A curve with equation $y = f(x)$ is such that $f'(x) = 6x^2 - \frac{8}{x^2}$. It is given that the curve passes through the point $(2, 7)$.

Find $f(x)$.

[3]

Cambridge Math & Physics

kambridgemp.com

- 2** The function f is defined by $f(x) = \frac{1}{3}(2x - 1)^{\frac{3}{2}} - 2x$ for $\frac{1}{2} < x < a$. It is given that f is a decreasing function.

Find the maximum possible value of the constant a .

[4]

Cambridge Math & Physics
kambridgemp.com

- 3** A line with equation $y = mx - 6$ is a tangent to the curve with equation $y = x^2 - 4x + 3$.

Find the possible values of the constant m , and the corresponding coordinates of the points at which the line touches the curve. [6]

Cambridge Math & Physics

kambridgemp.com

- 4 (a) Show that the equation

$$\frac{\tan x + \sin x}{\tan x - \sin x} = k,$$

where k is a constant, may be expressed as

$$\frac{1 + \cos x}{1 - \cos x} = k. \quad [2]$$

.....

.....

.....

.....

.....

.....

.....

- (b) Hence express $\cos x$ in terms of k . [2]

.....

.....

.....

.....

.....

.....

.....

- (c) Hence solve the equation $\frac{\tan x + \sin x}{\tan x - \sin x} = 4$ for $-\pi < x < \pi$. [2]

.....

.....


.....

.....

.....

.....

.....



- Math & Physics

- ambridge
kambrius

$$g(x) = x^2 + 4x + 13.$$

- Cambridge Math & Physics
kambridgemp.com

- [illegible]

- 7 (a) Write down the first four terms of the expansion, in ascending powers of x , of $(a - x)^6$. [2]

[illegible]

- (b) Given that the coefficient of x^2 in the expansion of $\left(1 + \frac{2}{ax}\right)(a - x)^6$ is -20 , find in exact form the possible values of the constant a . [5]

Cambridge Math & Physics

$$g : x \mapsto \frac{1}{2x+1} \text{ for } x < -\frac{1}{2}.$$

[4]

Cambridge Math & Physics

kanbridgemp.com

(b) Find an expression for $(fg)^{-1}(x)$.

[3]

Cambridge Math & Physics

kanbridgemp.com

- 9 (a)** A geometric progression is such that the second term is equal to 24% of the sum to infinity.

Find the possible values of the common ratio.

[3]

Cambridge Math & Physics

kambridgemp.com

- (b) An arithmetic progression P has first term a and common difference d . An arithmetic progression Q has first term $2(a + 1)$ and common difference $(d + 1)$. It is given that

$$\frac{\text{5th term of } P}{\text{12th term of } Q} = \frac{1}{3} \quad \text{and} \quad \frac{\text{Sum of first 5 terms of } P}{\text{Sum of first 5 terms of } Q} = \frac{2}{3}.$$

Find the value of a and the value of d .

[6]

- 10** Points $A(-2, 3)$, $B(3, 0)$ and $C(6, 5)$ lie on the circumference of a circle with centre D .

(a) Show that angle $ABC = 90^\circ$. [2]

(b) Hence state the coordinates of D . [1]

.....

.....


.....

.....

(c) Find an equation of the circle. [2]

ambridge
kambinus

- Cambridge Math & Physics
- kambridgemp.com



The diagram shows part of the curve with equation $y = x^{\frac{1}{2}} + k^2x^{-\frac{1}{2}}$, where k is a positive constant.

- (a) Find the coordinates of the minimum point of the curve, giving your answer in terms of k . [4]

Cambridge Math & English
kamdbridgemp.com

The tangent at the point on the curve where $x = 4k^2$ intersects the y-axis at P .

- (b) Find the y-coordinate of P in terms of k . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The shaded region is bounded by the curve, the x-axis and the lines $x = \frac{9}{4}k^2$ and $x = 4k^2$.

- (c) Find the area of the shaded region in terms of k . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Kambridge Math & Physics
kambridgeemp.com

Kambridge Math & Physics
kambridgemp.com

Kambridge Math & Physics
kambridgemp.com

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.