

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/13

Paper 1 Pure Mathematics 1 (P1)

October/November 2019

1 hour 45 minutes

Candidates answer on the Question Paper.

Additional Materials: List of Formulae (MF9)

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name in the spaces at the top of this page.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** the questions in the space provided. If additional space is required, you should use the lined page at the end of this booklet. The question number(s) must be clearly shown.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 75.

This document consists of **20** printed pages.

- 1** (i) Expand $(1 + y)^6$ in ascending powers of y as far as the term in y^2 . [1]

[illegible]

- (ii) In the expansion of $(1 + (px - 2x^2))^6$ the coefficient of x^2 is 48. Find the value of the positive constant p . [3]

Cambridge Math & Physics

- 2** The function g is defined by $g(x) = x^2 - 6x + 7$ for $x > 4$. By first completing the square, find an expression for $g^{-1}(x)$ and state the domain of g^{-1} . [5]


Cambridge Math & Physics

kambridgemp.com

- 3** The equation of a curve is $y = x^3 + x^2 - 8x + 7$. The curve has no stationary points in the interval $a < x < b$. Find the least possible value of a and the greatest possible value of b . [4]


Cambridge Math & Physics

kambridgemp.com



(i) Express angle CAO in radians in terms of π . [1]

(ii) Find the area of the shaded region in terms of r , π and $\sqrt{3}$, simplifying your answer. [4]



The dimensions of a cuboid are x cm, $2x$ cm and $4x$ cm, as shown in the diagram.

- (i) Show that the surface area S cm² and the volume V cm³ are connected by the relation

$$S = 7V^{\frac{2}{3}}.$$

[3]

Kambridge Math & Physics
kambridgemp.com

- (ii) When the volume of the cuboid is 1000 cm^3 the surface area is increasing at $2 \text{ cm}^2 \text{ s}^{-1}$. Find the rate of increase of the volume at this instant. [4]

Cambridge Math & Physics
kcambridgemp.com

- 6** A line has equation $y = 3kx - 2k$ and a curve has equation $y = x^2 - kx + 2$, where k is a constant.

(i) Find the set of values of k for which the line and curve meet at two distinct points. [4]

Cambridge Math & Physics

kambridgemp.com

- (ii) For each of two particular values of k , the line is a tangent to the curve. Show that these two tangents meet on the x -axis. [3]

Cambridge Math & Physics
kamdridgemp.com

- 7 (i) Show that the equation $3 \cos^4 \theta + 4 \sin^2 \theta - 3 = 0$ can be expressed as $3x^2 - 4x + 1 = 0$, where $x = \cos^2 \theta$. [2]

Cambridge Math & Physics
kamdbridgemp.com

Cambridge Math & Physics

kanbridgemp.com

- 8** A function f is defined for $x > \frac{1}{2}$ and is such that $f'(x) = 3(2x - 1)^{\frac{1}{2}} - 6$.

(i) Find the set of values of x for which f is decreasing.

[4]

Cambridge Math & Physics

kanbridgemp.com

(ii) It is now given that $f(1) = -3$. Find $f(x)$.

[4]

Kambridge Math & Physics
kambridgep.com

- 9 The first, second and third terms of a geometric progression are $3k$, $5k - 6$ and $6k - 4$, respectively.

(i) Show that k satisfies the equation $7k^2 - 48k + 36 = 0$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(ii) Find, showing all necessary working, the exact values of the common ratio corresponding to each of the possible values of k . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (iii) One of these ratios gives a progression which is convergent. Find the sum to infinity. [2]

$$\overrightarrow{OA} = \begin{pmatrix} -8 \\ -4 \\ 2 \end{pmatrix}, \quad \overrightarrow{OB} = \begin{pmatrix} 10 \\ 2 \\ 11 \end{pmatrix} \quad \text{and} \quad \overrightarrow{OX} = \begin{pmatrix} -2 \\ -2 \\ 5 \end{pmatrix}.$$

- [3]

Cambridge Math & Physics
kambriqemp.com

The position vector of a point C is given by $\overrightarrow{OC} = \begin{pmatrix} 1 \\ -8 \\ 3 \end{pmatrix}$.

(ii) Show that CX is perpendicular to AX .

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(iii) Find the area of triangle ABC .

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



- [illegible]

- (ii) Find, showing all necessary working, the volume of revolution obtained when the shaded region is rotated through 360° about the x -axis. [8]

Cambridge Math & Physics

kambridgemp.com

Cambridge Math & Physics

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.